

A NOTE ON A RESULT OF ZHANG ABOUT MONOCHROMATIC SUMSETS OF REALS

CHRIS LAMBIE-HANSON

ABSTRACT. We give an application of a higher-dimensional Δ -system lemma by using it in a slight modification of the proof of a recent result of Zhang about additive partition relations on the reals. This is meant to illustrate the use of the Δ -system lemma in question, and gives a slight improvement to the local version of Zhang's result.

The purpose of this note, which is not intended for publication, is to provide an exposition of a proof of a recent result of Zhang [4] in which a certain higher-dimensional Δ -system lemma used in [4] is replaced by a different higher-dimensional Δ -system lemma proven in [3]. Both lemmas involve starting with a sequence $\langle u_a \mid a \in [\mu]^n \rangle$ of sets of ordinals indexed by n -tuples from some cardinal μ , and then finding a set $H \subseteq \mu$ of some specified size such that $\langle u_a \mid a \in [H]^n \rangle$ satisfies certain uniformities. The advantage of our lemma in [3] is that, at least in the context of accessible cardinals, weaker assumptions are placed on the size of μ necessary to guarantee the existence of such a set H .

Zhang's result deals with partition relations for the additive structure $(\mathbb{R}, +)$. Given an additive structure $(A, +)$ and cardinals κ, r , the partition relation $A \rightarrow^+(\kappa)_r$ is the assertion that, for every coloring $c : A \rightarrow r$, there is $X \in [A]^\kappa$ such that $c \upharpoonright (X + X)$ is constant, where $X + X = \{x + y \mid x, y \in X\}$ (i.e., repetitions are allowed). Hindman, Leader, and Strauss prove in [1] that, if $2^{\aleph_0} < \aleph_\omega$, then there is $r < \omega$ such that $\mathbb{R} \not\rightarrow^+(\aleph_0)_r$. It was then shown by Komjáth et al. [2] that, modulo a large cardinal assumption, it is consistent that $\mathbb{R} \rightarrow^+(\aleph_0)_r$ for all $r < \omega$. This was improved by Zhang [4], who removed the large cardinal assumption and proved the following theorem.

Theorem 1 (Zhang, [4]). *Suppose that $\mathbb{P} = \text{Add}(\omega, \beth_\omega)$ is the forcing notion to add \beth_ω -many Cohen reals. Then in $V^{\mathbb{P}}$, we have $\mathbb{R} \rightarrow^+(\aleph_0)_r$ for all $r < \omega$.*

This shows that the result of Hindman, Leader, and Strauss is at least consistently sharp in the sense that, applying Zhang's result to a model of GCH, we obtain a forcing extension in which $2^{\aleph_0} = \aleph_{\omega+1}$ and $\mathbb{R} \rightarrow^+(\aleph_0)_r$ holds for all $r < \omega$.

An examination of Zhang's proof and the assumptions on the size of μ needed to prove the relevant Δ -system lemma shows that, for a fixed $r < \omega$, if \mathbb{P} is the forcing to add at least \beth_{4r}^+ -many Cohen reals, then $\mathbb{R} \rightarrow^+(\aleph_0)_r$ holds in $V^{\mathbb{P}}$. Our proof lowers this \beth_{4r}^+ to \beth_{2r}^+ , thus providing a local improvement to Zhang's result. We first make a note of some of our notational conventions.

Notation 2. If X is a set and κ is a cardinal, then $[X]^\kappa = \{Y \subseteq X \mid |Y| = \kappa\}$. If a is a set of ordinals, then $\text{otp}(a)$ denotes the order type of a under the natural ordering of the ordinals. We will frequently conflate sets of ordinals with increasing sequences of ordinals. So, for instance, if a is a set of ordinals and $i < \text{otp}(a)$,

then $a(i)$ is the unique $\eta \in a$ such that $\text{otp}(a \cap \eta) = i$. If $\mathbf{m} \subseteq \text{otp}(a)$, then $a[\mathbf{m}] = \{a(i) \mid i \in \mathbf{m}\}$. If a and b are sets of ordinals, then we write $a < b$ to mean that $\alpha < \beta$ for all $(\alpha, \beta) \in a \times b$.

A cardinal λ is said to be $<\kappa$ -inaccessible if $\nu^{<\kappa} < \lambda$ for all $\nu < \lambda$.

The proof presented here is essentially the same as that in [4]; we provide details just to verify that our Δ -system lemma is sufficient to carry out the proof. We first give two definitions from [2] and [4].

Definition 3. Suppose that μ is a cardinal, $m < \omega$, $a \in [\mu]^m$, and $s : m \rightarrow \mathbb{N}$. Then $s * a$ is the function from μ to \mathbb{N} defined by letting $s(a(i)) = s(i)$ for all $i < m$ and $s(\alpha) = 0$ for all $\alpha \in \mu \setminus a$. Notice that $s * a$ is then a member of $\bigoplus_{\alpha < \mu} \mathbb{N}$.

Definition 4. Suppose that $\ell \leq r < \omega$ and $2 \leq r$. Define a function $s_\ell^r : r + \ell \rightarrow \mathbb{N}$ by setting, for all $j < r + \ell$,

$$s_\ell^r(j) = \begin{cases} 2 & \text{if } j < 2\ell \\ 4 & \text{otherwise.} \end{cases}$$

We also need to recall some notation and results about higher-dimensional Δ -systems from [3].

Definition 5. Suppose that $\ell \leq r < \omega$, with $2 \leq r$. Then define a set $\mathbf{m}_\ell^r \subseteq 2r$ by letting $\mathbf{m}_\ell^r = \{2k + 1 \mid k < r\} \cup \{2k \mid k < \ell\}$. Notice that $|\mathbf{m}_\ell^r| = r + \ell$. Given a set $a \in [\text{On}]^{2r}$, let $a_\ell^r = a[\mathbf{m}_\ell^r]$.

Definition 6. Suppose that a and b are sets of ordinals.

- (1) We say that a and b are *aligned* if $\text{otp}(a) = \text{otp}(b)$ and, for all $\gamma \in a \cap b$, we have $\text{otp}(a \cap \gamma) = \text{otp}(b \cap \gamma)$.
- (2) If a and b are aligned then we let $\mathbf{r}(a, b) := \{i < \text{otp}(a) \mid a(i) = b(i)\}$. Notice that, in this case, $a \cap b = a[\mathbf{r}(a, b)] = b[\mathbf{r}(a, b)]$.

Definition 7. Suppose that H is a set of ordinals, $0 < n < \omega$, and, for all $b \in [H]^n$, u_b is a set of ordinals. We call $\langle u_b \mid b \in [H]^n \rangle$ a *uniform n -dimensional Δ -system* if there is an ordinal ρ and, for each $\mathbf{m} \subseteq n$, a set $\mathbf{r}_\mathbf{m} \subseteq \rho$ satisfying the following statements.

- (1) $\text{otp}(u_b) = \rho$ for all $b \in [H]^n$.
- (2) For all $a, b \in [H]^n$, if a and b are aligned, then u_a and u_b are aligned and, if $\mathbf{r}(a, b) = \mathbf{m}$, then $\mathbf{r}(u_a, u_b) = \mathbf{r}_\mathbf{m}$.
- (3) For all $\mathbf{m}_0, \mathbf{m}_1 \subseteq n$, we have $\mathbf{r}_{\mathbf{m}_0 \cap \mathbf{m}_1} = \mathbf{r}_{\mathbf{m}_0} \cap \mathbf{r}_{\mathbf{m}_1}$.

Definition 8. Suppose that $i < \rho$ are ordinals and $a, b \in [\text{On}]^\rho$. We say that a and b are *aligned above i* if $a[\rho \setminus i]$ and $b[\rho \setminus i]$ are aligned.

Definition 9. Suppose that a and b are sets of ordinals. Then the *intersection type of a and b* , denoted $\text{tp}_{\text{int}}(a, b)$, is the set $\{(i, j) \in \text{otp}(a) \times \text{otp}(b) \mid a(i) = b(j)\}$.

Definition 10. Suppose that I is a set and, for all $i \in I$, u_i is a set of ordinals. Then $\text{tp}(\langle u_i \mid i \in I \rangle)$ is a function from $\text{otp}(\bigcup_{i \in I} u_i)$ to $\mathcal{P}(I)$ defined as follows. First, let $\bigcup_{i \in I} u_i$ be enumerated in increasing order as $\langle \alpha_\eta \mid \eta < \text{otp}(\bigcup_{i \in I} u_i) \rangle$. Then, for all $\eta < \text{otp}(\bigcup_{i \in I} u_i)$, let $\text{tp}(\langle u_i \mid i \in I \rangle)(\eta) := \{i \in I \mid \alpha_\eta \in u_i\}$.

Intuitively, $\text{tp}(\langle u_i \mid i \in I \rangle)$ completely describes the order relations that hold between entries in $\langle u_i \mid i \in I \rangle$. We will often slightly abuse notation and write, for instance, $\text{tp}(u_0, u_1, u_2)$ instead of $\text{tp}(\langle u_0, u_1, u_2 \rangle)$.

Definition 11. Suppose that a is a nonempty set of ordinals and $i < \text{otp}(a)$.

- (1) We say that an ordinal α is i -possible for a if the following two statements hold:
 - (a) if $i > 0$, then $\alpha > a(i-1)$;
 - (b) if $i+1 < \text{otp}(a)$, then $\alpha < a(i+1)$.
Intuitively, α is i -possible for a if $a(i)$ can be replaced by α without changing the relative positions of the other elements of a .
- (2) If α is i -possible for a , then $a_{i \rightarrow \alpha}$ is the set $(a \setminus \{a(i)\}) \cup \{\alpha\}$, i.e., the set obtained by replacing the i^{th} element of a with α .

Definition 12. Given a regular cardinal λ , recursively define $\sigma(\lambda, n)$ for $1 \leq n < \omega$ by letting $\sigma(\lambda, 1) = \lambda$ and, given $1 \leq n < \omega$, letting $\sigma(\lambda, n+1) = (2^{<\sigma(\lambda, n)})^+$. Note that $\sigma(\lambda, n)$ is regular for each $1 \leq n < \omega$.

The following result is the higher-dimensional Δ -system lemma from [3].

Theorem 13. *Suppose that*

- $1 \leq n < \omega$;
- $\kappa < \lambda$ are infinite cardinals, λ is regular and $<\kappa$ -inaccessible, and $\mu = \sigma(\lambda, n)$;
- $g: [\mu]^n \rightarrow 2^{<\kappa}$;
- for all $b \in [\mu]^n$, we are given a set $u_b \in [\text{On}]^{<\kappa}$.

Then there are $H \subseteq \mu$ and $k < 2^{<\kappa}$ such that

- (1) $|H| = \lambda$;
- (2) $g(b) = k$ for all $b \in [H]^n$;
- (3) $\langle u_b \mid b \in [H]^n \rangle$ is a uniform n -dimensional Δ -system.

Moreover, if $n \geq 2$, for all $a, b \in [H]^n$ and all $k < n$, if it is the case that a and b are aligned above k and $a(k) = b(k)$, then, for any ordinal $\alpha \in H$ that is k -possible for both a and b , we have $\text{tp}_{\text{int}}(u_a, u_b) = \text{tp}_{\text{int}}(u_{a_{k \rightarrow \alpha}}, u_{b_{k \rightarrow \alpha}})$.

We are now ready to prove our adaptation of Zhang's result. As mentioned above, it is essentially the same as the proof from [4]. It was proven in [4] that $\mathbb{R} \rightarrow^+ (\aleph_0)_2$ holds in ZFC, so we only consider the case $r > 2$.

Theorem 14. *Suppose that $2 < r < \omega$ and \mathbb{P} is the forcing to add at least \beth_{2r}^+ -many Cohen reals. Then, in $V^{\mathbb{P}}$, we have $\mathbb{R} \rightarrow^+ (\aleph_0)_r$.*

Proof. Let $\mu = (\beth_{2r}^+)^V$, and let $\theta \geq \mu$ be a cardinal such that $\mathbb{P} = \text{Add}(\omega, \theta)$. We think of conditions of \mathbb{P} as being finite partial functions from θ to 2, ordered by reverse inclusion.

We identify $(\mathbb{R}, +)$ with $(\bigoplus_{\alpha < 2^\omega} \mathbb{Q}, +)$. We will actually show that, in $V^{\mathbb{P}}$, we have $\bigoplus_{\alpha < \mu} \mathbb{N} \rightarrow^+ (\aleph_0)_r$. Since we have $2^\omega \geq \theta \geq \mu$ in $V^{\mathbb{P}}$, this suffices.

Fix a \mathbb{P} -name \dot{c} for a function from $\bigoplus_{\alpha < \mu} \mathbb{N}$ to ω . We claim that the empty condition forces the existence of an infinite X such that $c \upharpoonright (X + X)$ is constant.

For each $\ell \leq r$, let \dot{d}_ℓ be a \mathbb{P} -name for the function from $[\mu]^{r+\ell}$ to r defined by letting $\dot{d}_\ell(a) = \dot{c}(s_\ell^r * a)$ for all $a \in [\mu]^{r+\ell}$.

For each $a \in [\mu]^{2r}$, let \mathcal{A}_a be a maximal antichain in \mathbb{P} such that, for each $q \in \mathcal{A}_a$ and each $\ell \leq r$, q decides the value of $\dot{d}_\ell(a_\ell^r)$. Since \mathbb{P} has the countable chain condition, each \mathcal{A}_a is countable, so we can enumerate it (possibly with repetitions) as $\langle q_{a,m} \mid m < \omega \rangle$. Let $u_{a,m} = \text{dom}(q_{a,m})$, and let $\bar{q}_{a,m} : \text{otp}(u_{a,m}) \rightarrow 2$ be

defined by letting $\bar{q}_{a,m}(i) = q_{a,m}(u_{a,m}(i))$ for all $i < \text{otp}(u_{a,m})$. For each $\ell \leq r$, let $w_{a,m,\ell} < r$ be such that $q_{a,m} \Vdash "d_\ell(a_\ell^r) = w_{a,m,\ell}"$. Let $u_a = \bigcup_{m < \omega} u_{a,m}$.

Now the map g that takes $a \in [\mu]^{2r}$ to the triple

$$\langle \langle \bar{q}_{a,m} \mid m < \omega \rangle, \langle w_{a,m,\ell} \mid m < \omega, \ell \leq r \rangle, \text{tp}(\langle u_a \rangle \frown \langle u_{a,m} \mid j < \omega \rangle) \rangle$$

can easily be coded as a map from $[\mu]^{2r}$ to $2^{<\omega_1}$. Moreover, u_a is countable for all $a \in [\mu]^{2r}$, and $\mu = \beth_{2r}^+ = \sigma(\beth_1^+, 2r)$. Since \beth_1^+ is $<\omega_1$ -inaccessible, we can apply Theorem 13 to $\langle u_a \mid a \in [\mu]^{2r} \rangle$ and g to obtain $H \subseteq \mu$ of size \beth_1^+ and a fixed triple $\tau = \langle \langle \bar{q}_m \mid m < \omega \rangle, \langle w_{m,\ell} \mid m < \omega, \ell \leq r \rangle, t \rangle$ such that $g(a) = \tau$ for all $a \in [H]^{2r}$ and $\langle u_a \mid a \in [H]^{2r} \rangle$ is a uniform $2r$ -dimensional Δ -system and satisfies the “moreover” clause in the statement of Theorem 13. Let $\rho < \omega_1$ be such that $\text{otp}(u_a) = \rho$ for all $a \in [H]^{2r}$, and let $\langle \mathbf{r}_m \subseteq \rho \mid \mathbf{m} \subseteq 2r \rangle$ witness that $\langle u_a \mid a \in [H]^{2r} \rangle$ is a uniform $2r$ -dimensional Δ -system.

Fix sets $\langle A_k \mid k < r \rangle$ such that each A_k is a subset of H of order type $\omega + 1$ and $A_k < A_{k'}$ for all $k < k' < r$. Let $\alpha_0^k = \min(A_k)$ and $\alpha_\omega^k = \max(A_k)$ for all $k < r$. We identify elements of $\prod_{k < r} [A_k]^2$ as elements of $[\mu]^{2r}$ in the obvious way.

Let G be \mathbb{P} -generic over V , and let c and $\langle d_\ell \mid \ell \leq r \rangle$ be the realizations of \dot{c} and $\langle \dot{d}_\ell \mid \ell \leq r \rangle$, respectively, in $V[G]$. For every $a \in [H]^{2r}$, there is a unique $m_a < \omega$ such that $q_{a,m_a} \in G$. Working now in $V[G]$, we will recursively construct a matrix of ordinals $\langle \alpha_j^k \mid k < r, j < \omega \rangle$ such that, for each $k < r$, $\langle \alpha_j^k \mid j < \omega \rangle$ is an increasing sequence of ordinals in $A_k \setminus \{\alpha_\omega^k\}$ (note that we have already specified $\alpha_0^k = \min(A_k)$). At the end, we will let $A_k^* = \{\alpha_j^k \mid j \leq \omega\}$. Our construction will be by recursion on the anti-lexicographic order on $r \times \omega$, i.e., we set $(k, j) < (k', j')$ iff $j < j'$ or $(j = j'$ and $k < k')$. To specify the requirements our construction will satisfy, we need some further definitions.

At the end of the construction, an element $a \in \prod_{k < r} [A_k^*]^2$ will be called *canonical* if $a = \{\alpha_{j_0}^0, \alpha_{j'_0}^0, \alpha_{j_1}^1, \alpha_{j'_1}^1, \dots, \alpha_{j_{r-1}}^{r-1}, \alpha_{j'_{r-1}}^{r-1}\}$, where

- for each $k < r$, we have $j_k < j'_k$;
- for each $k_0 < k_1 < r$, we have $j_{k_0} < j_{k_1}$;
- for each $k < r$, we have $j_{r-1} < j'_k$;
- for each $k_0 < k_1 < r$, if $j'_{k_0} < \omega$, then $j'_{k_0} \leq j'_{k_1}$.

If $a = \{\alpha_{j_0}^0, \alpha_{j'_0}^0, \dots, \alpha_{j_{r-1}}^{r-1}, \alpha_{j'_{r-1}}^{r-1}\} \in \prod_{k < r} [A_k^*]^2$ is canonical, then the *index* of a is the set $\{j_k \mid k < r\}$. Note that this is an element of $[\omega]^r$. In our construction, we will arrange so that, for every canonical $a \in \prod_{k < r} [A_k^*]^2$ and every $\ell \leq r$, the value of $d_\ell(a_\ell^r)$ depends only on the index of a . This will be arranged in the following way: for each canonical $a = \{\alpha_{j_0}^0, \alpha_{j'_0}^0, \dots, \alpha_{j_{r-1}}^{r-1}, \alpha_{j'_{r-1}}^{r-1}\} \in \prod_{k < r} [A_k^*]^2$, let $\hat{a} = \{\alpha_{j_0}^0, \alpha_\omega^0, \dots, \alpha_{j_{r-1}}^{r-1}, \alpha_\omega^{r-1}\}$. In other words, \hat{a} is the canonical element of $\prod_{k < r} [A_k^*]^2$ with the same index as a and whose other elements are precisely the elements of $\{\alpha_\omega^k \mid k < r\}$. We will ensure that, for every canonical element a , we have $m_a = m_{\hat{a}}$. It will follow that $d_\ell(a_\ell^r) = d_\ell(\hat{a}_\ell^r) = w_{m_{\hat{a}}, \ell}$.

We now describe the construction of $\langle \alpha_{k,j} \mid k < r, j < \omega \rangle$. We have already specified $\alpha_{k,0}$ for all $k < r$. Now fix $(k^*, j^*) \in r \times \omega$ with $j^* \geq 1$, and suppose that we have defined $\alpha_{k,j}$ for all $(k, j) < (k^*, j^*)$. For each $k < r$, let $B_k = \{\alpha_{k,j} \mid (k, j) < (k^*, j^*)\} \cup \{\alpha_{k,\omega}\}$, i.e., B_k is the portion of A_k^* that has already been specified. The notion of a canonical element of $\prod_{k < r} [B_k]^2$ is straightforwardly inherited from the

notion of a canonical element of $\prod_{k < r} [A_k^*]^2$. Our recursion hypothesis is simply that, for every canonical element a , we have $m_a = m_{\hat{a}}$.

We call a canonical element $a = \{\alpha_{j_0}^0, \alpha_{j'_0}^0, \dots, \alpha_{j_{r-1}}^{r-1}, \alpha_{j'_{r-1}}^{r-1}\}$ of $\prod_{k < r} [B_k]^2$ relevant if $j'_k = \omega$ for all k with $k^* \leq k < r$. Let

$$q^* = \bigcup \{q_{a, m_a} \mid a \text{ is a relevant canonical element}\}.$$

Since there are only finitely many relevant canonical elements, we have $q^* \in G$. Also, for each relevant canonical element a and each $\alpha \in A_{k^*} \setminus (\{\alpha_\omega^{k^*}\} \cup \alpha_{j_{k^*}^{k^*}})$, let $a_\alpha = a_{(2k^*+1) \rightarrow \alpha} = (a \setminus \{\alpha_\omega^{k^*}\}) \cup \{\alpha\}$.

Claim 15. *There is $\alpha \in A_{k^*} \setminus (\{\alpha_\omega^{k^*}\} \cup \alpha_{j_{k^*}^{k^*}})$ such that, for every relevant canonical element a , we have $m_{a_\alpha} = m_a$, i.e., $q_{a_\alpha, m_a} \in G$.*

Proof. Assume not. Note that, since there are only finitely many canonical relevant elements, each of which is a finite set of ordinals and hence in V , the statement of the claim is expressible in V as a statement in the forcing language for \mathbb{P} . Therefore, since the claim fails, we can fix a single condition $s \in G$ that forces its failure. Assume without loss of generality that $s \leq q^*$.

Let $\mathbf{m} = 2r \setminus \{2k^* + 1\}$, and let $C = A_{k^*} \setminus (\{\alpha_\omega^{k^*}\} \cup \alpha_{j_{k^*}^{k^*}})$. For each relevant canonical element a , the set $\{u_{a_\alpha} \mid \alpha \in C\}$ is a Δ -system whose root is equal to $u_{a_\alpha}[\mathbf{r}_\mathbf{m}]$ for each $\alpha \in C$. Since there are only finitely many relevant canonical elements a and since $\text{dom}(s)$ is finite, we can therefore fix $\alpha \in C$ such that, for every relevant canonical element a , we have $(u_{a_\alpha} \setminus u_{a_\alpha}[\mathbf{r}_\mathbf{m}]) \cap \text{dom}(s) = \emptyset$. Let

$$q^{**} = s \cup \bigcup \{q_{a_\alpha, m_a} \mid a \text{ is a relevant canonical element}\}.$$

We claim that q^{**} is a condition in \mathbb{P} , i.e., it is actually a function. To see this, it suffices to verify the following two statements:

- For every relevant canonical element a , we have $s \parallel q_{a_\alpha, m_a}$.
- For every pair of relevant canonical elements a and b , we have $q_{a_\alpha, m_a} \parallel q_{b_\alpha, m_b}$.

To verify the first statement, fix a relevant canonical element a . By our choice of α , we have $\text{dom}(q_{a_\alpha, m_a}) \cap \text{dom}(s) \subseteq u_{a_\alpha}[\mathbf{r}_\mathbf{m}]$. But a_α and a are aligned, with $\mathbf{r}(a_\alpha, a) = \mathbf{m}$, so $u_{a_\alpha}[\mathbf{r}_\mathbf{m}] = u_a[\mathbf{r}_\mathbf{m}]$. By the fact that g is constant on $[H]^{2r}$, we have $q_{a_\alpha, m_a} \upharpoonright u_{a_\alpha}[\mathbf{r}_\mathbf{m}] = q_{a, m_a} \upharpoonright u_a[\mathbf{r}_\mathbf{m}]$. But $s \leq q_{a, m_a}$, so $s \leq q_{a_\alpha, m_a} \upharpoonright u_{a_\alpha}[\mathbf{r}_\mathbf{m}]$, so $s \parallel q_{a_\alpha, m_a}$.

To verify the second statement, fix a pair of relevant canonical elements, a and b . It easily follows from the definitions of “relevant” and “canonical” that a and b are aligned above $2k^* + 1$. Moreover, we have $a(2k^* + 1) = b(2k^* + 1) = \alpha_\omega^{k^*}$. Therefore, by the “moreover” clause of Theorem 13, we have $\text{tp}_{\text{int}}(u_a, u_b) = \text{tp}_{\text{int}}(u_{a_\alpha}, u_{b_\alpha})$. Now suppose for sake of contradiction that $q_{a_\alpha, m_a} \perp q_{b_\alpha, m_b}$. Then there is $\gamma \in \text{dom}(q_{a_\alpha, m_a}) \cap \text{dom}(q_{b_\alpha, m_b})$ such that $q_{a_\alpha, m_a}(\gamma) \neq q_{b_\alpha, m_b}(\gamma)$. Fix $i_a, i_b < \rho$ such that $\gamma = u_{a_\alpha}(i_a) = u_{b_\alpha}(i_b)$. Then $(i_a, i_b) \in \text{tp}_{\text{int}}(u_{a_\alpha}, u_{b_\alpha})$, so $(i_a, i_b) \in \text{tp}_{\text{int}}(u_a, u_b)$, so there is δ such that $\delta = u_a(i_a) = u_b(i_b)$. By the fact that g is constant on $[H]^{2r}$, we have

$$q_{a, m_a}(\delta) = q_{a_\alpha, m_a}(\gamma) \neq q_{b_\alpha, m_b}(\gamma) = q_{b, m_b}(\delta),$$

and hence $q_{a, m_a} \perp q_{b, m_b}$. But, by assumption, we have $q_{a, m_a}, q_{b, m_b} \in G$, which is a contradiction.

This finishes the verification that q^{**} is a condition. But now note that q^{**} extends s and forces that α witnesses the truth of the claim, contradicting our choice of s . Therefore, the claim holds. \square

We can now let $\alpha_{j^*}^{k^*}$ be any α witnessing the truth of Claim 15. Let us verify that this maintains the recursion hypothesis. For $k < r$, let $B'_k = B_k$ if $k \neq k^*$, and let $B'_{k^*} = B_{k^*} \cup \{\alpha_{j^*}^{k^*}\}$. Fix a canonical element a of $\prod_{k < r} [B'_k]^2$. We must show that $m_a = m_{\hat{a}}$. By the recursion hypothesis, we may assume that $\alpha_{j^*}^{k^*} \in a$. Note that, for all $k < r$ with $k > k^*$, we have not yet defined $\alpha_{j^*}^k$. Therefore, by the definition of “canonical element”, we must be in one of two cases:

Case 1: $\alpha_{j^*}^{k^*} = a[2k^*]$ and $k^* = r - 1$. Again by the definition of “canonical element”, it must be the case here that $a[2k + 1] = \alpha_{\omega}^k$ for all $k < r$. Hence, $a = \hat{a}$, so the recursion hypothesis is trivially satisfied.

Case 2: $\alpha_{j^*}^{k^*} = a[2k^* + 1]$. Here, it must be the case that $a[2k + 1] = \alpha_{\omega}^k$ for all $k < r$ with $k > k^*$. Let $b = a_{(2k^*+1) \rightarrow \alpha_{\omega}^{k^*}}$, and, for notational simplicity, let $\alpha = \alpha_{j^*}^{k^*}$. Then b is a relevant canonical element of $\prod_{k < r} [B_k]^2$. Notice that $a = b_{\alpha}$, so by our choice of α , we have $m_a = m_b$. By our recursion hypothesis, we have $m_b = m_{\hat{b}}$. But $\hat{b} = \hat{a}$, so $m_a = m_{\hat{a}}$.

We have thus maintained our recursion hypothesis and can move on to the next step of the construction. This therefore completes our construction of $\langle A_k^* \mid k < r \rangle$.

The rest of the proof is exactly as in [4], but we provide a sketch for completeness. By our construction of $\langle A_k^* \mid k < r \rangle$, for each $\ell \leq r$ we have a well defined function $f_{\ell} : [\omega]^r \rightarrow r$ such that, for each $y \in [\omega]^r$ and each canonical $a \in \prod_{k < r} A_k^*$, if the index of a is y , then $d_{\ell}(a_{\ell}^r) = f_{\ell}(y)$. By Ramsey’s theorem, there is an infinite $Y \subseteq \omega$ such that each f_{ℓ} is constant on $[Y]^r$, say with value $\varepsilon_{\ell} < r$. By throwing away the elements of $\omega \setminus Y$ and reindexing, we may assume for notational simplicity that $Y = \omega$, i.e., for every canonical $a \in \prod_{k < r} A_k^*$ and every $\ell \leq r$, we have $d_{\ell}(a_{\ell}^r) = \varepsilon_{\ell}$.

By the pigeonhole principle, there are $\ell_0 < \ell_1 \leq r$ such that $\varepsilon_{\ell_0} = \varepsilon_{\ell_1} =: \varepsilon$. For all $j < \omega$, define $a_j \in \prod_{k < \ell_0} [A_k^*]^2 \times \prod_{\ell_0 \leq k < r} A_k^*$ by specifying that a_j contains the following:

- $\{\alpha_k^k, \alpha_{\omega}^k\}$ for each $k < \ell_0$;
- $\{\alpha_{k+(j+1)r}^k\}$ for $\ell_0 \leq k < \ell_1$;
- $\{\alpha_{\omega}^k\}$ for each $\ell_1 \leq k < r$.

Note that $a_j \in [H]^{r+\ell_0}$. Let $x_j = \frac{1}{2} s_{\ell_0}^r * a_j \in \bigoplus_{\alpha < \mu} \mathbb{N}$, and let $X = \{x_j \mid j < \omega\}$. We claim that $c \upharpoonright (X + X)$ is constant with value ε . There are two things to verify.

First, we must show that $c(x_j + x_j) = \varepsilon$ for all $j < \omega$. Thus, fix $j < \omega$. Let $a = a_j \cup \{a_k^k \mid \ell_0 \leq k < r\}$. Then a is a canonical element of $\prod_{k < r} [A_k^*]^2$ and $a_{\ell_0}^r = a_j$. Therefore, we have

$$c(x_j + x_j) = c(s_{\ell_0}^r * a_j) = d_{\ell_0}(a_{\ell_0}^r) = \varepsilon_{\ell_0} = \varepsilon,$$

as desired.

Next, we must show that $c(x_j + x_{j'}) = \varepsilon$ for all $j < j' < \omega$. Thus, fix $j < j' < \omega$. Let $a = a_j \cup a_{j'} \cup \{a_k^k \mid \ell_1 \leq k < r\}$. The following facts are easily verified.

- a is a canonical element of $\prod_{k < r} [A_k^*]^2$.
- $a_{\ell_1}^r = a_j \cup a_{j'}$.
- $x_j + x_{j'} = s_{\ell_1}^r * (a_j \cup a_{j'})$.

As a result, we have

$$c(x_j + x_{j'}) = c(s_{\ell_1}^r * (a_j \cup a_{j'})) = d_{\ell_1}(a_{\ell_1}^r) = \varepsilon_{\ell_1} = \varepsilon.$$

We have thus shown that $c \upharpoonright (X + X)$ is constant with value ε , thus finishing the proof. \square

REFERENCES

1. Neil Hindman, Imre Leader, and Dona Strauss, *Pairwise sums in colourings of the reals*, Abh. Math. Semin. Univ. Hambg. **87** (2017), no. 2, 275–287. MR 3696151
2. Péter Komjáth, Imre Leader, Paul A. Russell, Saharon Shelah, Dániel T. Soukup, and Zoltán Vidnyánszky, *Infinite monochromatic sumsets for colourings of the reals*, Proc. Amer. Math. Soc. **147** (2019), no. 6, 2673–2684. MR 3951442
3. Chris Lambie-Hanson, *Higher-dimensional Delta-systems*, (2020), Preprint.
4. Jing Zhang, *Monochromatic sumset without large cardinals*, Fund. Math. (2020), To appear.

INSTITUTE OF MATHEMATICS OF THE CZECH ACADEMY OF SCIENCES, ŽITNÁ 25, PRAHA 1, CZECHIA

Email address: `lambiehanson@math.cas.cz`

URL: `http://math.cas.cz/lambiehanson`