

MAZUR INTERSECTION PROPERTY FOR ASPLUND SPACES

MIROSLAV BAČÁK AND PETR HÁJEK

ABSTRACT. The main result of the present note states that it is consistent with the ZFC axioms of set theory (relying on Martin's Maximum MM axiom), that every Asplund space of density character ω_1 has a renorming with the Mazur intersection property. Combined with the previous result of Jiménez and Moreno (based upon the work of Kunen under the continuum hypothesis) we obtain that the MIP renormability of Asplund spaces of density ω_1 is undecidable in ZFC.

The Mazur intersection property (MIP for short) was first investigated by S. Mazur in [10] as a purely geometrical isometric property of a Banach space, and has since been studied extensively over the years. An early result of Mazur claims that a Banach space with a Fréchet differentiable norm (necessarily an Asplund space) has the MIP ([10]). Phelps [12] proved that a separable Banach space has a MIP renorming if and only if its dual is separable, or equivalently, if it is an Asplund space. Much of the further progress in the theory depended on an important characterization of MIP, due to Giles, Gregory and Sims [6], by the property that w^* -denting points of B_{X^*} are norm dense in S_{X^*} . This result again suggests a close connection of MIP to Asplund spaces, as the latter can be characterized in a similar way as spaces such that bounded subsets of their dual are w^* -dentable. It has opened a way to applying biorthogonal systems to the MIP. Namely, Jiménez and Moreno [9] have proved that if a Banach space X^* admits a fundamental biorthogonal system $\{(x_\alpha, f_\alpha)\}$, where f_α belong to $X \subset X^{**}$, then X has a MIP renorming. As a corollary to this criterion ([9]), they get that every Banach space can be embedded into a Banach space which is MIP renormable, a rather surprising result which in particular strongly shows that MIP and Asplund properties, although closely

Date: March 2008.

2000 Mathematics Subject Classification. 46B03.

Key words and phrases. Mazur intersection property, Asplund space, Martin's Maximum axiom, fundamental biorthogonal system.

Supported by grants: Institutional Research Plan AV0Z10190503, GAČR 201/07/0388, GA UK 72 407, GA ČR 201/07/0394.

related, are distinct. We remark that MIP has connections with other parts of Banach space theory, such as the ball topology ([7]) etc..

On the other hand, all known examples of an Asplund space without MIP renorming ([9], [8]), such as $C(K)$ space, where K is a compact space constructed by Kunen ([11]), were constructed using additional set theoretical assumptions, such as the continuum hypothesis CH (or \clubsuit). In fact $C(K)$, where K is Kunen's compact, has density character ω_1 , the smallest density for which such a result may hold in view of Phelps' theorem.

The main result of the present note goes in the positive direction. Namely, it is consistent with the ZFC axioms of set theory (relying on a powerful additional axiom - Martin's Maximum MM [5]), that every Asplund space of density character ω_1 has a renorming with the Mazur intersection property. This answers a question of Godefroy, communicated to the authors by Zizler. We thank V. Zizler for suggesting the problem and sharing his insights with us. Combined with the previous result of Jiménez and Moreno we obtain that the MIP renormability of Asplund spaces of density ω_1 is undecidable in ZFC.

Our proof has three main ingredients. It combines a recent result of Todorčević, claiming that under MM every Banach space with density ω_1 has a fundamental system, with the machinery of projectional generators, and a criterion of MIP renormability due to Jiménez and Moreno via fundamental systems. We recall that MM is an axiom consistent with ZFC, see [5] for details. We prefer not to state MM explicitly, as we do not enter into any axiomatic arguments here, and our reliance on MM is solely through the use of Todorčević's theorem in [13] (see also [8]). In order to explain in some detail the role of these results, we need some preliminary definitions.

Let $(X, \|\cdot\|)$ be a Banach space, denote its closed unit ball B_X and its unit sphere S_X . More generally, $B(x, \rho) = \{y \in X : \|y - x\| \leq \rho\}$ is a ball centered at x with radius $\rho \geq 0$. The dual of X is denoted X^* , and by w^* we mean the weak* topology on X^* , that is $\sigma(X^*, X)$. The symbol τ_p stands for the topology of pointwise convergence in $c_0(\omega_1)$. The set of rational numbers and the linear hull with rational coefficients are denoted \mathbb{Q} , \mathbb{Q} -span, respectively.

A Banach space X is an *Asplund space* if every separable subspace of X has a separable dual space. We say that a Banach space $(X, \|\cdot\|)$ has the *Mazur intersection property* if every bounded closed convex set K is an intersection of closed balls. More precisely, $K = \bigcap_\alpha B(x_\alpha, \rho_\alpha)$, for some system of points $x_\alpha \in X$, and radii ρ_α .

Let X be a Banach space and Γ a nonempty set. A family $\{(x_\gamma, x_\gamma^*)\}_{\gamma \in \Gamma}$ of pairs $x_\gamma \in X$, $x_\gamma^* \in X^*$ is called a *biorthogonal system* in $X \times X^*$ if $x_\alpha^*(x_\beta) =$

$\delta_{\alpha\beta}$, where $\delta_{\alpha\beta}$ is Kronecker's delta, for all $\alpha, \beta \in \Gamma$. A biorthogonal system is *fundamental* if $\overline{\text{span}}\{x_\gamma\}_{\gamma \in \Gamma} = X$.

If A is a subset of a Banach space X and $B \subset X^*$, we say that A *countably supports* B , (or that B is *countably supported* by A), provided $\{a \in A : b(a) \neq 0\}$ is at most countable, for all $b \in B$. A subset $N \subset X^*$ is *1-norming* if $\|x\| = \sup\{x^*(x) : x^* \in N \cap B_{X^*}\}$, for all $x \in X$.

For a nonseparable Banach space X of density μ , we define *projectional resolution of identity (PRI)* as a family $\{P_\alpha : \omega \leq \alpha \leq \mu\}$ of linear projections on X such that $P_\omega = 0$, P_μ is the identity mapping on X , and for all $\omega \leq \alpha, \beta \leq \mu$:

- (i) $\|P_\alpha\| = 1$,
- (ii) $\text{dens}P_\alpha X \leq \text{card}\alpha$,
- (iii) $P_\alpha P_\beta = P_\beta P_\alpha = P_\alpha$, if $\alpha \leq \beta$,
- (iv) $\cup_{\alpha < \beta} P_{\alpha+1} X$ is norm dense in $P_\beta X$, if $\omega < \beta$.

A standard way of obtaining PRI on Banach spaces is using the Projectional Generator. This technique has an advantage of allowing some additional properties for the PRI, important in our proof.

Let X be a Banach space and $W \subset X^*$ be 1-norming set closed under linear combinations with rational coefficients. Let $\Phi : W \rightarrow 2^X$ be an at most countably valued mapping such that for every nonempty set $B \subset W$ with linear closure, $\Phi(B)^\perp \cap \overline{B}^{w^*} = \{0\}$. Then the couple (W, Φ) is called a *projectional generator (PG)*.

Let $\{P_\alpha : \omega \leq \alpha \leq \mu\}$ be a PRI on X , $G \subset X$. We will say that the given PRI is *subordinated* to the set G if $P_\alpha(x) \in \{0, x\}$ for all α and $x \in G$.

We are now going to present the main theorems, whose combination leads to the proof of our main result.

Theorem 1. [8, Theorem 3.42]

Let X be a nonseparable Banach space with a projectional generator (W, Φ) , and a set $G \subset X$ that countably supports W . Then X has a PRI $\{P_\alpha : \omega \leq \alpha \leq \mu\}$, which is subordinated to the set G .

We shall use the following theorem of Jiménez and Moreno ([9], see also [8, Theorem 8.42]).

Theorem 2. *Let X^* be a dual Banach space with a fundamental biorthogonal system $\{(x_\alpha^*, x_\alpha)\}_{\alpha < \omega_1} \subset X^* \times X^{**}$, with the property that $x_\alpha \in X \subset X^{**}$. Then X admits an equivalent norm with the Mazur intersection property.*

The next theorem, due to Todorčević ([13], see also [8, Theorem 4.48]), is the main ingredient of our proof.

Theorem 3. (MM) *Every Banach space of density ω_1 has a fundamental biorthogonal system.*

For proofs of the above results, additional references and more information we refer to the recent book [8]. The main result of our note is following theorem.

Theorem 4. (MM) *Let X be an Asplund space of density ω_1 . Then X admits an equivalent norm with the Mazur intersection property.*

Proof. Because X is an Asplund space, we have that $\text{dens}X^* = \omega_1$. (One way of seeing this is to use the Jayne-Rogers continuous selectors of the duality mapping together with the Bishop-Phelps' theorem, see [2] for details). According to Theorem 2, it suffices to find a fundamental biorthogonal system of X^* , with $\{(x_\alpha^*, x_\alpha)\}_{\alpha < \omega_1} \subset X^* \times X$. We claim that if there exists a linear continuous mapping $T : (X^*, w^*) \rightarrow (c_0(\omega_1), \tau_p)$ with a nonseparable range, such that

$$(0.1) \quad T(x^*) = (x^*(y_\gamma))_{\gamma \in \Gamma}, \quad \text{for some } \Gamma \text{ uncountable, and } y_\gamma \in S_X,$$

then X^* has a fundamental biorthogonal system $\{(x_\alpha^*, x_\alpha)\}_{\alpha < \omega_1} \subset X^* \times X$. The proof of this claim follows by inspection of Todorčević's argument in [13] (or [8, Theorem 4.45]) which shows the same implication without w^* -topology involved. As a matter of fact, during the construction of a fundamental system, one only passes to a suitable long subsequence of $\{y_\gamma\}$, which then becomes the $\{x_\alpha\} \subset X \subset X^{**}$.

It remains to find $\{y_\gamma\}_{\gamma \in \Gamma} \subset S_X$, Γ uncountable, that define the operator in (0.1). By Theorem 3, there exists a fundamental biorthogonal system in X , denoted by $\{(e_\gamma, f_\gamma)\}_{\gamma < \omega_1} \subset X \times X^*$. We may without loss of generality assume that $\|e_\gamma\| = 1$ for all $\gamma < \omega_1$. We shall show that the desired sequence $\{y_\gamma\}_{\gamma \in \Gamma}$ can be found as an uncountable subsequence of $(e_\gamma)_{\gamma < \omega_1}$, (hence $\Gamma \subset \omega_1$).

Since X is an Asplund space, there exists a projectional generator (W, Φ) on X^* , where $W = \text{span}\{e_\gamma\}_{\gamma < \omega_1}$. Indeed, the set W is dense in X , and thus it is 1-norming for X^* . By [3, Proposition 8.2.1], there exists a projectional generator (X, Φ) on X^* , so the restriction of Φ to W will give the needed mapping. Note that the set $G = \{f_\gamma\}_{\gamma < \omega_1}$ is subordinated to (W, Φ) . By Theorem 1, there exists a PRI $\{P_\alpha : \omega \leq \alpha \leq \omega_1\}$ and such that

$$(0.2) \quad P_\alpha(f_\gamma) = \begin{cases} f_\gamma, & \text{or} \\ 0, & \end{cases}$$

for all $\omega \leq \alpha \leq \omega_1$ and $\gamma \in \omega_1$. We claim that we are allowed to put another additional condition on our PRI, namely we require that

$$(0.3) \quad P_\alpha^*(e_\gamma) = \begin{cases} e_\gamma, & \text{or} \\ 0, & \end{cases}$$

and moreover $P_\alpha^*(e_\gamma) = e_\gamma$ if and only if $P_\alpha(f_\gamma) = f_\gamma$.

This is achieved by entering into the construction of PRI using the PG (W, Φ) . In fact, reading through the proof of Theorem 3.42 in [8], we see that the auxiliary multi-valued mapping $\Psi : X \rightarrow 2^W$ may be without loss of generality chosen so that $e_\gamma \in \Psi(f_\gamma)$ for all $\gamma < \omega_1$. We also adjust the original Φ (which in the case of an Asplund space may be chosen a sequence of Jayne-Rogers' selectors restricted to W) by adding finitely many extra elements $\{f_{\gamma_i}\}_{i=1}^n$ to the set $\Phi(v)$, for every $v = \sum_{i=1}^n r_i e_{\gamma_i}$, $r_i \in \mathbb{R}$. The adjusted pair (W, Φ) is again a PG for X^* . In the case of the first adjustment it is obvious, because the defining properties of the originally chosen Ψ have not been violated. In the second case, it suffices to note a simple general fact, that for any given PG (W, Φ) , upon replacing Φ by any countably valued mapping $\tilde{\Phi}$, with the property that $\Phi(f) \subset \tilde{\Phi}(f)$, the pair $(W, \tilde{\Phi})$ will remain a PG. This is apparent from the definition of PG, since $\tilde{\Phi}(B) \supseteq \Phi(B)$ implies $(\tilde{\Phi}(B))^\perp \subseteq \Phi(B)^\perp$. Having these extra conditions at hand, it is easy to observe that the set B_α in the proof of Theorem 3.42 in [8] equals to the \mathbb{Q} -span of the (countable) set $\{e_\lambda\}_{\lambda \in \Lambda_\alpha}$, where $\Lambda_\alpha = \{\gamma : e_\gamma \in B_\alpha\}$. This implies, using Goldstine's theorem, that $P_\alpha^*(X^{**}) = \overline{\text{span}}^{w^*} \{e_\lambda : \lambda \in \Lambda_\alpha\}$ and also $(\text{Id} - P_\alpha^*)(X^{**}) = \overline{\text{span}}^{w^*} \{e_\lambda : \lambda \notin \Lambda_\alpha\}$. Thus $P_\alpha^*(e_\gamma) = e_\gamma$ if $\gamma \in \Lambda_\alpha$ and $P_\alpha^*(e_\gamma) = 0$ otherwise. Denote $\Gamma_\alpha = \{\gamma < \omega_1 : f_\gamma \in (P_{\alpha+1} - P_\alpha)X^*\}$. Let $\Gamma = \{\alpha : \Gamma_\alpha \neq \emptyset\}$. We claim that $\text{card}\Gamma = \omega_1$. Indeed, $P_\alpha(X^*)$ is a separable Banach space for every $\alpha < \omega_1$, while $\text{span}\{f_\gamma\}_{\gamma < \omega_1}$ is nonseparable. For convenience we may without loss of generality assume that $\Gamma = [\omega_1, \omega_1)$. For every $\alpha \in \Gamma$ we choose $\gamma(\alpha) < \omega_1$, such that $\gamma(\alpha) \in \Gamma_\alpha$. Since the mapping

$$x^* \mapsto (\|(P_{\alpha+1} - P_\alpha)(x^*)\|)_{\omega \leq \alpha < \omega_1}$$

is into $c_0([\omega, \omega_1))$ (Lemma VI.1.2 in [2]), we have that the mapping

$$T : x^* \mapsto (e_{\gamma(\alpha)}((P_{\alpha+1} - P_\alpha)(x^*)))_{\alpha \in \Gamma}$$

is into $c_0(\Gamma)$. However, it is a consequence of (0.3) that $e_{\gamma(\alpha)} = e_{\gamma(\alpha)} \circ (P_{\alpha+1} - P_\alpha)$ on X^* . It follows that

$$T : x^* \mapsto (x^*(e_\gamma))_{\gamma \in \Gamma}$$

is a $w^* - \tau_p$ continuous operator from X^* into $c_0(\Gamma)$. Further, from (0.2) and the choice of Γ , we get that the range of T is not separable. This finishes the proof. \square

We will conclude the paper by making a few remarks. Note that on the one hand, the dual to our original Asplund space, X^* , admits a PG, so it has an M-basis, and thus it has a bounded injection into $c_0(\omega_1)$ (for these facts see [8]). On the other hand, X^* in general does not admit injections into $c_0(\omega_1)$, that are continuous in the $w^* - \tau_p$ topology, since the last condition characterizes WCG spaces X , and not every Asplund space of density ω_1 is WCG. The main point of our construction is that under MM one can actually construct a "large" (nonseparable range) operator from X^* into $c_0(\omega_1)$, which is at the same time $w^* - \tau_p$ continuous.

There remains a number of interesting questions on MIP in Asplund spaces. An example of an Asplund space with MIP but having no Fréchet renorming is not known. (According to a result of Deville [1], the long James space has MIP renorming, but the existence of Fréchet renorming on this space is open.) It is an open problem due to Zizler, if an Asplund space with a C^1 -smooth bump function is MIP renormable. It seems to be open whether the statement claiming that every Asplund space is MIP renormable, is consistent in ZFC.

REFERENCES

- [1] R. Deville: *Un Théoreme de transfert pour la propriété des boules*. Canad. Math. Bull. 30 (1987), 295–300.
- [2] R. Deville, G. Godefroy and V. Zizler: *Smoothness and renormings in Banach spaces*. Monographs and Surveys in Pure and Applied Mathematics 64, Pitman, 1993.
- [3] M.J. Fabian: *Gâteaux differentiability of convex functions and topology*. John Wiley & Sons, Inc., New York, 1997.
- [4] M. Fabian, P. Habala, P. Hájek, V. Montesinos, J. Pelant and V. Zizler: *Functional analysis and infinite dimensional geometry*. Canadian Math. Soc. Books (Springer-Verlag), 2001.
- [5] M. Foreman, M. Magidor and S. Shelah: *Martin's Maximum, saturated ideals, and non-regular ultrafilters. Part I*. Ann. Math. 127 (1988), 1–47.
- [6] J.R. Giles, D.A. Gregory and B. Sims: *Characterization of normed linear spaces with Mazur's intersection property*. Bull. Austral. Math. Soc. 18 (1978), 471–476.
- [7] G. Godefroy and N.J. Kalton: *The ball topology and its applications*. Contemp. Math. 85 (1989), 195–237.
- [8] P. Hájek, V. Montesinos, J. Vanderwerff, V. Zizler: *Biorthogonal systems in Banach spaces*. Springer, New York, 2008.
- [9] M. Jiménez-Sevilla and J.P. Moreno: *Renorming Banach Spaces with the Mazur Intersection Property*. J. Funct. Anal. 144 (1997), 486–504.
- [10] S. Mazur: *Über schwache Konvergenz in en Raumen L^p* . Studia Math. 4 (1933), 128–133.
- [11] S. Negrepontis: *Banach Spaces and Topology*. Handbook of set-theoretic topology, 1045–1142, North-Holland, Amsterdam, 1984.
- [12] R.R. Phelps: *A representation theorem for bounded convex sets*. Proc. AMS 11 (1960), 976–983.

- [13] S. Todorčević: *Biorthogonal systems and quotient spaces via Baire category theorem*.
Math. Ann. 335 (2006), 687–715.

DEPARTMENT OF MATHEMATICAL ANALYSIS, CHARLES UNIVERSITY, SOKOLOVSKÁ 83,
186 75 PRAHA 8, CZECH REPUBLIC

E-mail address: `bacak@karlin.mff.cuni.cz`

MATHEMATICAL INSTITUTE, CZECH ACADEMY OF SCIENCE, ŽITNÁ 25, 115 67 PRAHA
1, CZECH REPUBLIC

E-mail address: `hajek@math.cas.cz`