

Exercises for Mathematical Logic (Fall 2022/23)

We have seen in the lecture that the De Morgan language $\{\wedge, \vee, \neg, \top, \perp\}$ is functionally complete (on $\{0, 1\}$), and specifically, that every Boolean function can be represented by a CNF or DNF of size $O(2^n)$.

1. Prove that $\{\vee, \neg\}$, $\{\rightarrow, \perp\}$, and $\{\uparrow\}$ are functionally complete, where $x \uparrow y$ denotes the Sheffer stroke $\neg(x \wedge y)$.

2. Prove that $\{\rightarrow\}$, $\{\wedge, \vee, \top, \perp\}$, and $\{\leftrightarrow, \top, \perp\}$ are not functionally complete.

[Hint: Find a nontrivial property of Boolean functions which is preserved by composition, and holds for functions in the given basis.]

3. For any Boolean function $f: \{0, 1\}^n \rightarrow \{0, 1\}$, the following are equivalent:

(i) $\{f\}$ is functionally complete.

(ii) $f(0, \dots, 0) = 1$, $f(1, \dots, 1) = 0$, and there exists an assignment $\langle e_0, \dots, e_{n-1} \rangle \in \{0, 1\}^n$ such that $f(e_0, \dots, e_{n-1}) = f(\neg e_0, \dots, \neg e_{n-1})$.

[Hint: For (ii) \rightarrow (i), look at functions obtained from f by identifying some of the variables.]

4. For any $n \in \mathbb{N}$, the *parity* function $\bigoplus_{i < n} x_i: \{0, 1\}^n \rightarrow \{0, 1\}$ is defined as $\sum_{i < n} x_i \pmod 2$. Show that any DNF or CNF representing $\bigoplus_{i < n} x_i$ has size $\Omega(2^n)$. [Hint: What terms of the form $\bigwedge_{i \in I} x_i^{e_i}$ can imply one of $\bigoplus_{i < n} x_i = 0$ or $\bigoplus_{i < n} x_i = 1$? Here, $I \subseteq [n]$, $e_i \in \{0, 1\}$, $x^1 = x$, $x^0 = \neg x$.]

5. There are formulas representing $\bigoplus_{i < n} x_i$ of size $O(n^c)$ for some constant c .

[Hint: Consider a balanced tree of binary parities. You may get it down to $c = 2$.]

6. Any DNF equivalent to the CNF $\bigwedge_{i < n} (x_i \vee y_i)$ has size $\Omega(2^n)$.

7. Prove the propositional soundness theorem: for all $\Gamma \subseteq \text{Prop}(A)$ and $\varphi \in \text{Prop}(A)$, if $\Gamma \vdash \varphi$, then $\Gamma \models \varphi$.

8. Let $\Gamma, \Delta \subseteq \text{Prop}(A)$ and $\varphi, \psi \in \text{Prop}(A)$. Show that if $\Gamma \vdash \varphi$ and $\Delta, \varphi \vdash \psi$, then $\Gamma, \Delta \vdash \psi$.

9. For every $\varphi \in \text{Prop}(A)$, we define its *De Morgan dual* $\varphi^d \in \text{Prop}(A)$ by induction on the complexity of φ :

$$\begin{aligned} a^d &= a, & a \in A, & & (\neg\varphi)^d &= \neg(\varphi^d), \\ \top^d &= \perp, & & & \perp^d &= \top, \\ (\varphi \wedge \psi)^d &= (\varphi^d \vee \psi^d), & & & (\varphi \vee \psi)^d &= (\varphi^d \wedge \psi^d). \end{aligned}$$

Show that for all assignments $v: A \rightarrow \{0, 1\}$, $v(\varphi^d) = v_{\neg}(\neg\varphi)$, where $v_{\neg}: A \rightarrow \{0, 1\}$ is the assignment defined by $v_{\neg}(a) = 1 - v(a)$ for each $a \in A$.

10. Let $\varphi, \psi \in \text{Prop}(A)$.

(i) $\varphi \equiv \psi$ if and only if $\varphi^d \equiv \psi^d$.

(ii) $\varphi \models \psi$ if and only if $\psi^d \models \varphi^d$.

11. (If you are familiar with topology.) Give a direct proof of the propositional compactness theorem, not using the completeness theorem.

[Hint: Consider the product topology on the set $\{0, 1\}^A$ of all assignments.]

12. Prove that if a term $t(x_0, \dots, x_{n-1})$ is free for y in a formula $\varphi(x_0, \dots, x_{n-1}, y)$, then for all terms s_0, \dots, s_{n-1} , the formula $(\varphi(t/y))(s_0/x_0, \dots, s_{n-1}/x_{n-1})$ is syntactically identical to the formula $\varphi(s_0/x_0, \dots, s_{n-1}/x_{n-1}, t(s_0/x_0, \dots, s_{n-1}/x_{n-1})/y)$.

13. A set of propositional or first-order sentences S is *independent* if S is not equivalent to S' for any proper subset $S' \subsetneq S$.

(i) S is independent iff $S \setminus \{\varphi\} \not\models \varphi$ for all $\varphi \in S$.

(ii) Show that every countable theory T has an independent axiomatization, i.e., an independent set of sentences S equivalent to T . [Hint: Try to generalize the fact that $\{\varphi, \psi\} \equiv \{\varphi, \psi \vee \neg\varphi\}$.]

(Uncountable theories have independent axiomatizations as well by a theorem of I. Reznikoff, but this is more difficult to prove.)

14. Consider a modification of the first-order proof system given in the lecture such that the axioms of equality are replaced with the axiom $x = x$ and the axiom schema $t = s \wedge \varphi(t/s) \rightarrow \varphi(s/x)$ for all formulas φ and terms t, s free for x in φ . Show that this is equivalent to the original proof system.

15. For any formula $\varphi(x)$ and variable y free for x in φ , show that the formula $\exists y (\exists x \varphi(x) \rightarrow \varphi(y))$ is provable.

16. Using Vaught's test, show the completeness of the theory of a successor: it has a language with one unary function symbol s , and axioms $s(x) = s(y) \rightarrow x = y$, $\forall x \exists y s(y) = x$, and $s^n(x) \neq x$ for each $n \in \mathbb{N}_{>0}$, where s^n denotes the n -fold iteration of s (i.e., $s^0(x)$ is x , and s^{n+1} is $s(s^n(x))$).

17. For each $n \in \mathbb{N}$, let P_n denote the path graph of length n , i.e., the structure $\langle [n], E_n \rangle$, where $[n] = \{0, \dots, n-1\}$ and $E_n = \{\langle i, j \rangle \in [n]^2 : |i-j| = 1\}$. Show that there is no sentence φ such that for all $n \in \mathbb{N}$, $P_n \models \varphi$ iff n is odd. [Hint: Adapt the previous exercise.]

18. Fix a field F . The theory of vector spaces over F has a language consisting of the language $\{+, -, 0\}$ of abelian groups and unary functions $a \cdot x$ for each $a \in F$; it has the usual algebraic axioms (axioms of abelian groups, $ab \cdot x = a \cdot (b \cdot x)$, $1 \cdot x = x$, $(a+b) \cdot x = a \cdot x + b \cdot x$, $a \cdot (x+y) = a \cdot x + a \cdot y$). Show that the theory of infinite vector spaces over F (i.e., with additional axioms $\exists x_0 \dots \exists x_n \bigwedge_{i < j} x_i \neq x_j$ for $n \in \mathbb{N}$) is complete and κ -categorical for all infinite $\kappa > |F|$. [Hint: Every vector space has a basis.]

19. Prove $\mathbb{Q} \vdash \forall x (x \leq \bar{n} \vee \bar{n} \leq x)$ for each $n \in \mathbb{N}$.

20. \mathbb{Q} proves $x \cdot y = 0 \rightarrow x = 0 \vee y = 0$, and more generally, $x \cdot y = \bar{n} \rightarrow x = 0 \vee y \leq \bar{n}$ for each $n \in \mathbb{N}$.

21. The standard model \mathbb{N} extends to an L_{PA} -structure \mathbb{N}^∞ with domain $\mathbb{N} \cup \{\infty\}$, $\infty \notin \mathbb{N}$, so that $\mathbb{N}^\infty \models \mathbb{Q}$. Moreover, we are free to choose $(0 \cdot \infty)^{\mathbb{N}^\infty}$ in an arbitrary way (while the rest of the model is uniquely determined by the axioms of \mathbb{Q}). Conclude that \mathbb{Q} does not prove any of the formulas $S(x) \not\leq x$, $x \cdot y = y \cdot x$, or $0 \cdot x \neq 1$.

22. \mathbb{Q} does not prove $x + y = y + x$ or $0 + (x + y) = (0 + x) + y$.

[Hint: Modify the previous exercise to a model with two "infinities".]

In the next three exercises, you will develop an alternative sequence encoding scheme due to Edward Nelson.

23. The set $\{x : \exists n \in \mathbb{N} x = 2^n\}$ of powers of 2 is definable by a Δ_0 formula, not using the 2^n function. [Hint: Consider the divisors of x .]

24. Consider an encoding of finite sets $X \subseteq \mathbb{N}$ by pairs $\langle r, w \rangle$ where the binary expansion of r acts as a "ruler" with marks at positions of 1s, and the binary expansion of w is a concatenation of binary expansions of elements of X such that each element occupies the position between two ruler marks. Show that the predicate " x is in the set encoded by $\langle r, w \rangle$ " is Δ_0 -definable.

25. Construct a Δ_0 encoding of finite sequences based on the previous exercise.

A function $f: \mathbb{N} \rightarrow \mathbb{N}$ is *represented* by a formula $\varphi(x, y)$ in a theory T if $T \vdash \forall y (\varphi(\bar{n}, y) \leftrightarrow y = \bar{m})$ for all $n, m \in \mathbb{N}$ such that $f(n) = m$.

26. All partial computable functions are representable in \mathbf{Q} . [Hint: Using Σ_1 -definability of the graph of f , adapt the witness comparison argument from the proof of representability of decidable sets.]

27. All Σ_1 -definable sets are semidecidable.

28. (Craig's trick.) Every semidecidable theory is recursively axiomatizable. [Hint: Express $\text{Thm}(T)$ as $\exists y P(x, y)$ with P decidable. Given $x = \ulcorner \varphi \urcorner$ and y , devise a sentence equivalent to φ that encodes y .]

29. Show that every decidable consistent theory T has a decidable completion. [Hint: Consider a completion procedure that enumerates sentences φ one by one, and extends the current list of axioms with φ or $\neg\varphi$, whichever maintains consistency with T .]

30. Prove Gödel's diagonal lemma: for every formula $\varphi(x)$, there exists a sentence α such that $\mathbf{Q} \vdash \alpha \leftrightarrow \varphi(\ulcorner \alpha \urcorner)$. [Hint: Using representability of a suitable computable function (see Exer. 26), construct a formula $\psi(x)$ such that $\mathbf{Q} \vdash \psi(\ulcorner \chi \urcorner) \leftrightarrow \varphi(\ulcorner \chi(\ulcorner \chi \urcorner) \urcorner)$ for all $\chi(x)$.]